

Tier levels determine how quickly a laboratory report must be entered into EpiTrax by BEPHI Surveillance staff



Tier 1 (1 hour - Notify Immediately)

Anthrax
Botulism
Cholera
Dengue Fever
Diphtheria
Hantavirus Pulmonary Syndrome
Haemophilus influenzae, invasive disease
H. Influenza (If under 5 years old)
Hepatitis A IgM+ ONLY
Hepatitis B, acute
Influenza deaths in children <18 years of age
Measles (rubeola)
Meningococcemia
Mumps
Pertussis (whooping cough)
Plague (Yersinia pestis)
Poliomyelitis
Q Fever (Coxiella burnetii)
Rabies, animal
Rabies, human
Rubella, including congenital rubella syndrome
Smallpox
Tularemia
Varicella
Viral hemorrhagic fever

Tier 2 (1 business day)

Amebiasis
Brucellosis
Campylobacter infections
Cryptosporidiosis
Cyclospora infection
Escherichia coli O157:H7 (and other shiga toxin-producing E. coli, also known as STEC)
Giardiasis
Listeriosis
Hemolytic uremic syndrome, postdiarrheal
Salmonellosis, including typhoid fever
Shigellosis
Trichinosis

**ALL KHEL LABS NO MATTER
WHAT DISEASE TIER**

Tier 3 (2 business days)

Arboviral disease
Babesiosis
Coccidioidomycosis
Ehrlichiosis
Hepatitis B during pregnancy
Hepatitis, viral (acute and chronic)
Lyme disease
Malaria
Meningitis, bacterial
Psittacosis
Spotted Fever Rickettsiosis (RMSF)
Spongiform encephalopathy (STE) or prion disease (includes vCJD)
Streptococcal invasive, drug-resistant disease from Group A Streptococcus or Streptococcus pneumoniae
Streptococcus pneumoniae, invasive <5 years of age
(*notify VPD Coordinator*)
Tetanus
Toxic shock syndrome, streptococcal and staphylococcal
Vibriosis (not cholera)
Yellow fever
Yersiniosis

As of August 2012

